

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Naturally Occurring Asbestiform Minerals in Italian Western Alps and in Other Italian Sites

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1715911> since 2020-02-26T18:14:05Z

Published version:

DOI:10.2113/EEG-2276

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Naturally Occurring Asbestiform Minerals in Italian Western Alps and in Other Italian Sites

BELLUSO Elena

*Department of Earth Sciences, University of Torino, via Valperga Caluso n.35, 10125 Torino, Italy;
00393358345402, elena.belluso@unito.it
Interdepartmental Centre “G. Scansetti” for Studies on Abestos and Other Toxic Particulates,
University of Torino, Italy
Institute of Geosciences and Earth Resources -IGG, National Research Council –CNR of Italy*

BARONNET Alain

*Centre Interdisciplinaire de Nanosciences de Marseille (CINaM), Université Aix-Marseille, France;
Campus de Luminy, Case 913, 13288 Marseille Cedex 9, 00336 62 90 38 06, baronnet@cinam.univ-
mrs.fr*

CAPELLA Silvana

*Department of Earth Sciences, University of Torino, via Valperga Caluso n.35, 10125 Torino, Italy;
0039116705121, silvana.capella@unito.it
Interdepartmental Centre “G. Scansetti” for Studies on Abestos and Other Toxic Particulates,
University of Torino, Italy*

Abstract

The natural occurrence of asbestos (NOA) in rocks and soil has been known for many years in several areas of the world, differently from the natural presence of asbestiform minerals. In Italy the mapping of NOA is mandatory according to the 2001 and 2003 regulations. The investigation, not yet concluded, has revealed that in Italy NOA are represented by chrysotile, tremolite asbestos with minor amount of actinolite asbestos and anthophyllite asbestos. A field survey carried out in Italian Western Alps (IWA), dealing with the naturally occurring of asbestiform minerals non-asbestos classified (NONA) and not regulated, started many years ago and is still ongoing. It revealed that the following kinds of asbestiform silicates are present (in decreasing order of frequency): asbestiform polygonal serpentine and asbestiform antigorite, asbestiform diopside, asbestiform carlosturanite, asbestiform

forsterite, asbestiform sepiolite, asbestiform balangeroite, and asbestiform talc. The asbestiform non-silicates brugnatellite and brucite have been rarely detected. Outside the IWA, asbestiform zeolite (erionite and offretite), asbestiform sodium amphibole (fluoro-edenite), and few other asbestiform silicates have been also detected. For some asbestiform minerals, the identification is problematic and needs the use of transmission electron microscope (TEM) combining either the imaging at high magnification as the electron diffraction and the chemical data. This investigation is particularly important to distinguish four kinds of asbestiform minerals (antigorite, polygonal serpentine, carlosturanite, and balangeroite) from chrysotile since only the last one is regulated. The issue is much more complicated by the intergrowth of different fibrous species on the scale of sub micrometre.

Keywords: NOA; naturally occurring non-asbestos classified asbestiform minerals (NONA); TEM-EDS identifications; fibre intergrowth

Introduction

In recent years and in several countries the health investigations dealing with asbestos moved from the occupational to environmental exposure (e.g., Baumann et al., 2015; Abakay et al., 2016; Noonan, 2017). According some authors in fact the exposure to low but continuous exposure, as in the case of inhabitants of houses next to the asbestos bearing rock outcrops (i.e. naturally occurring asbestos: NOA), could cause health problems (e.g., Luce et al., 2000; Bernardini et al., 2003). The same problem could concern the exposure to naturally occurring non-asbestos classified asbestiform minerals (NONA). For some of them, the carcinogenicity to high dose exposure is known as in the case of asbestiform fluoeredenite, asbestiform erionite, asbestiform winchite, and asbestiform richterite (e.g., Burrigato et al., 2005, IARC, 2012). In addition to these last minerals, many others having an asbestiform morphology have been discovered over the years in different part of the world. The most

55 striking current example is represented by asbestiform antigorite. For many years this antigorite
56 asbestiform variety has not been recognized as having a full identity and little information has appeared
57 in few publications (e.g., Keeling et al., 2006).

58 But, exactly as it happens for the NOA, also the NONA can be dispersed in air both for natural
59 causes (weathering, natural atmospheric agents, landslide) and for anthropogenic causes (e.g.
60 excavation works).

61 If these asbestiform minerals are noxious following continuous low dose and/or sporadic high
62 doses, it will be known only in many years, as it has happened for example for asbestos given the long
63 latency time of related asbestos pathologies. For the instance, it would be a good practice to map the
64 presence of these minerals in every country.

65 As it concerns Italy, the mapping of NOA is mandatory according to the March 23, 2001 Law, n.
66 93 and to the related March 18, 2003, Environment Ministry Decree, n. 101. The mapping of NOA
67 must be carried out by each single Region through its own Environmental Protection Agency, using
68 literature data, geological maps, historical information on research permits and mining concessions,
69 reports and environmental monitoring carried out as part of environmental and strategic impact
70 assessment procedures for the construction of infrastructural works, analytical certificates, mining
71 activities of lithotypes suspected for the presence of asbestos, surveys for the Geological Cartography
72 project at 1: 50,000 scale, specific investigations and surveys, possible inspections and sometimes
73 collection of samples and laboratory analyses. Each year the individual EPA send any possible
74 additional data to the Ministry of Health that integrates it to the aim of the evaluation of the necessary
75 remediation works.

76 Currently, the map of the Italian NOA is published from the Ministry of the Environment and
77 Protection of the Territory and the Sea; it is updated to the year 2018 and reports NOA sites only from
78 three regions (Asbestos Mapping, 2018) but to the current state of scientific knowledge, NOA are

present in at least 7 other regions. This paper presents the different NOA and NONA until now detected in Italy.

Detected NOA and NONA

Firstly, owing to the different used definitions, it needs to define the meaning of the used words. According to the dimensional definition of the World Health Organization (WHO, 1997), many regulatory agencies and the literature (e.g. Belluso et al., 2017), in this note we use the definition listed below.

Fiber: inorganic particle with length $\geq 5 \mu\text{m}$, width $\leq 3 \mu\text{m}$, length/width (aspect ratio) $\geq 3:1$, parallel sides when seen in two dimensions, perpendicularly to fiber axis.

Fibril: a single mineral fiber which cannot be further separated longitudinally into smaller components (without losing the fibrous properties or appearances).

Asbestiform: adjective for fibers non asbestos classified having the “fiber” dimensions and at least one of the asbestos properties as flexibility, splitting etc.

Fiber bundle: parallel aggregate of mineral fibers.

In the scientific literature many researches touch the NOA topic. Some of them have been carried out specifically for mineralogical interests, others concern petrological, structural engineering or geology studies; sometimes the scientific investigations arise from minerals collectors. In any case, their identification is carried out by using different techniques, in some cases by using two or more than two in a complementary way. The most diffused identification techniques are X-ray powder diffractometry (XRPD), infrared spectroscopy (with Fourier transform), optical (OM) and electron microscopy (EM), usually scanning (SEM) and uncommonly transmission (TEM), coupled with energy dispersive spectrometry (EDS).

102 Based on scientific literature data, NOA are widespread in many areas of Italy and represented by
103 chrysotile and tremolite asbestos (both very diffused), the less diffused actinolite asbestos and finally
104 anthophyllite asbestos that is the much less diffused than others (e.g., Cavallo and Rimoldi, 2013;
105 Gaggero et al., 2013; Vignaroli et al., 2013; Gaggero et al., 2017). The Figure 1 shows the 10 Regions
106 where NOA are present in rocks, at the current state of knowledge. Crocidolite and amosite do not
107 constitute NOA in Italy.

108 Unlike NOA, the mapping of NONA is not mandatory in Italy. Several spot investigations,
109 carried out during both the asbestos investigation and specific mineralogical researches, show that
110 NONA are present in many Italian areas (8 Italian Regions) and abundant in some places. Except
111 Piedmont and Aosta Valley Regions which we will discuss later, the NONA identified in Italy are the
112 following (in alphabetic order): asbestiform antigorite, asbestiform Ca-erionite (with Ca-levyne),
113 asbestiform F-edinite, asbestiform gedrite, asbestiform Mg-horneblende, asbestiform offretite,
114 asbestiform polygonal serpentine, asbestiform sepiolite (e.g., Cattaneo et al., 2011; Bloise et al.,
115 2014;.Bloise et al., 2016; Bloise et al., 2017; Giordani et al., 2017; Lucci et al., 2018; Mattioli et al.,
116 2018) The Italian Regions where these asbestiform minerals have been detected are shown in Figure 2.

117 Regarding the Italian Western Alps (IWA) Regions, i.e. Piedmont and Aosta Valley, a field
118 survey dealing with NOA is carried out since 1980 and it is still ongoing (Baronnet and Belluso, 2002;
119 Belluso et al., 2015; Leone, 2018; Paccagnella, 2018). More than 300 samples have been collected and
120 analyzed by using XRPD coupled with SEM-EDS and/or TEM-EDS. As NOA, chrysotile, asbestos
121 tremolite, and, in lesser amount, asbestos actinolite have been detected.

122 The investigation in this area revealed that not only NOA are present, but that there are many
123 kinds of asbestiform silicates (sometimes in very high amount) of NONA, not regulated in Italy. Ten
124 are the identified NONA and eight of these are magnesium-containing silicates (Table 1). The complete
125 list is shown below, in decreasing order of frequency: asbestiform polygonal serpentine and

126 asbestiform antigorite (with very high frequency); asbestiform diopside, asbestiform carlosturanite,
127 asbestiform forsterite, asbestiform sepiolite, asbestiform balangeroite, and asbestiform talc. Rare are
128 asbestiform brugnatellite and asbestiform brucite, the only two non-silicates asbestiform minerals.

129 As far as quantities are concerned, asbestiform carlosturanite and asbestiform balangeroite are
130 very abundant. As regards the quantities of asbestiform antigorite and asbestiform polygonal
131 serpentine, is not possible to define the amount because these two minerals are always intergrown with
132 others fibrous minerals, asbestos and non asbestos classified.

133 For some asbestiform minerals, the certain identification is not so easy because they show similar
134 (at times even equal) characteristics on a macroscopic scale. As it concerns NOA and NONA in IWA
135 Regions, on the macroscopic scale in most cases it is not possible distinguish among chrysotile,
136 asbestiform antigorite, asbestiform polygonal serpentine, asbestiform carlosturanite, and asbestiform
137 balangeroite. Figure 3 shows an example of this similarity on the eye scale, both in outcropping rocks
138 and after picking fibers from the rock, between asbestiform antigorite (Figs. 3a and 3b) and chrysotile
139 (Fig. 3c). The bundles of fibres are flexible and their characteristics are very similar to each other. Also
140 the colour is similar because chrysotile containing a little amount of iron has not white colour but
141 cream to light brown.

142 Often it needs the use of the transmission electron microscope (TEM) combining either the
143 imaging at high magnification as the selected area electron diffraction (SAED) and the chemical data
144 (EDS).

145 This investigation is particularly important to distinguish four kinds of NONA and specifically
146 asbestiform antigorite (Fig. 4), asbestiform polygonal serpentine (Fig. 4), asbestiform carlosturanite,
147 and asbestiform balangeroite, from chrysotile (Fig. 4) since only the last one is regulated, being
148 asbestos classified. The issue is very complicated because usually, on the scale of little thousands or

149 even hundreds micrometers, the fibers of two to four mineral species are intergrown (owing to solid
150 state transformations) as it can see in Fig. 4.

151 Based on the detailed investigation by TEM-EDS, practically each bundle of fibers turns out to be
152 made up of at least 2 fibrous mineral species, except in the case of asbestiform sepiolite, which on the
153 other hand appears to have only intergrown with an organic phase (aliphatic hydrocarbons: Giustetto et
154 al., 2014).

155

156 **Conclusions**

157 In Italy NOA and NONA are diffused in several areas. NOA in Italy are represented only by four
158 mineral species; among them only chrysotile and tremolite asbestos are abundant and diffused whereas
159 actinolite asbestos is less diffused and anthophyllite asbestos is rare. NONA in Italy are constituted by
160 15 mineral species, many more than NOA, although they have not the same diffusion and abundance.

161 Detailed investigation on NOA and NONA, even if not completed, were conducted only in
162 Piedmont and in Aosta Valley Regions and lead to reveal many different NONA in rocks and in the
163 same bundle of fibers. Owing to the intergrowth on the sub-micrometric scale of different fibrous
164 species, many times only the TEM-EDS investigation allows to detect all the mineral species and for
165 certain phases only this kind of techniques allows to obtain a certain identification, as for example to
166 distinguish among asbestiform antigorite, asbestiform polygonal serpentine, and chrysotile, therefore
167 from non-asbestos and asbestos classified minerals.

168 Finally, many issues arise from the shown data as the relevance of the correct and representative
169 sampling, the importance of the use of the technique suitable to identify the content of these composite
170 materials (i.e. the bundle of fibers), the need of the extended surveys for extensive investigations where
171 there are rocks that may contain mineral fibers.

172 The investigation by TEM-EDS of fiber bundles collected around the world may give surprises
173 like those from Italian Western Alps rocks.

174

- 176 Abakay, A.; Tanrikulu, A. C. ; Ayhan, M. ; Imamoglu, M. S. ; Taylan, M. ; Kaplan, M. A.; and
177 Abakay, O., 2016, High-risk mesothelioma relation to meteorological and geological condition and
178 distance from naturally occurring asbestos: *Environmental Health Preventive Medicine*, Vol. 21, pp.
179 82–90.
- 180 Baronnet, A.; and Belluso, E., 2002, Microstructures of the silicates: key information about mineral
181 reactions and a link with the Earth and materials sciences: *Mineralogical Magazine*, Vol. 66, pp. 709-
182 735.
- 183 Baumann, F.; Buck, B.J.; Metcalf, R.V.; McLaurin, B.T.; Merkler, D.J.; and Carbone, M., 2015, The
184 presence of asbestos in the natural environment is likely related to mesothelioma in young individuals
185 and women from Southern Nevada: *Journal Thoracic Oncology*, Vol. 10, pp. 731–737.
- 186 Belluso, E.; Cavallo, A.; and Halterman, D., 2017, Crystal habit of mineral fibres: *European*
187 *Mineralogical Union Notes in Mineralogy*, Vol. 18, pp. 65-109.
- 188 Belluso, E.; Compagnoni, R.; and Ferraris, G., 1995, Occurrence of asbestiform minerals in the
189 serpentinites of the Piemonte Zone, Western Alps. In Politecnico di Torino (Editor), *Giornata di Studio*
190 *in ricordo del Prof. Stefano Zucchetti*, Torino, pp. 57-64.
- 191 Bernardini, P.; Schettino, B.; Sperduto, B.; Giannandrea, F.; Burragato, F.; and Castellino, N., 2003,
192 Three cases of pleural mesothelioma and environmental pollution with tremolite outcrops in Lucania:
193 *Giornale Italiano Medicina Lavoro Ergonomia*, Vol. 25, pp. 408–411.
- 194 Bloise, A.; Catalano, M.; Critelli, T.; Apollaro, C.; and Miriello, D., 2017, Naturally occurring
195 asbestos: potential for human exposure, San Severino Lucano (Basilicata, Southern Italy):
196 *Environmental Earth Sciences*, Vol. 76, pp. 648-660.
- 197 Bloise, A.; Critelli, T.; Catalano, M.; Apollaro, C.; Miriello, D.; Croce, A.; Barrese, E.; Liberi, F.;
198 Piluso, E.; Rinaudo, C.; and Belluso, E., 2014, Asbestos and other fibrous minerals contained in the

199 serpentinites of the Gimigliano-Mount Reventino Unit (Calabria, S-Italy): *Environmental Earth*
200 *Sciences*, Vol. 71, pp. 3773–3786.

201 Bloise, A.; Punturo, R.; Catalano, M.; Miriello, D.; and Cirrincione, R., 2016, Naturally occurring
202 asbestos (NOA) in rock and soil and relation with human activities: the monitoring example: *Italian*
203 *Journal Geosciences*, Vol. 135, pp. 268-279.

204 Burragato, F.; Comba, P.; Baiocchi, V.; Palladino, D.M.; Simeì, S.; Gianfagna, A.; Paoletti, L.; and
205 Pasero, R., 2005, Geo-volcanological, mineralogical and environmental aspects of quarry materials
206 related to pleural neoplasm in the area of Biancavilla, Mount Etna (Eastern Sicily, Italy):
207 *Environmental Geology*, Vol. 47, pp. 855-868.

208 Cattaneo, A.; Rossotti, A.; Pasquaré, G.; Somigliana, A.; and Cavallo, D. M., 2011, Analysis of fibrous
209 zeolites in the volcanic deposits of the Viterbo Province, Italy: *Environmental Earth Sciences*, Vol. 63,
210 pp. 861–871.

211 Cavallo, A.; and Rimoldi, B., 2013, Chrysotile asbestos in serpentinite quarries: A case study in
212 valmalenco, central alps, northern Italy: *Environmental Sciences: Processes Impacts*, Vol. 15, pp.
213 1341-1350.

214 Gaggero, L.; Crispini, L.; Isola, E.; and Marescotti, P., 2013, Asbestos in natural and anthropic
215 ophiolitic environments: a case study of geohazards related to the Northern Apennine ophiolites
216 (Eastern Liguria, Italy): *Ofioliti*, Vol. 38, pp. 29-40.

217 Gaggero, L.; Sanguineti, E.; Gonzalez, A. Y.; Militello, G. M.; Scuderi, A.; and Parisi G., 2017,
218 Airborne asbestos fibres monitoring in tunnel excavation: *Journal of Environmental Management*, Vol.
219 196, pp. 583-593.

220 Giordani, M.; Mattioli, M.; Ballirano, P.; Pacella, A.; Cenni, M.; Boscardin, M.; and Valentini, L.,
221 2017, Geological occurrence, mineralogical characterization, and risk assessment of potentially

222 carcinogenic erionite in Italy: *Journal Toxicology Environmental Health, Part B, Critical Reviews*,
 223 Vol. 20, pp. 81–103.

224 Giustetto, R.; Seenivasan, K.; and Belluso, E., 2014, Asbestiform sepiolite coated by aliphatic
 225 hydrocarbons from Perleto, Aosta Valley Region (Western Alps, Italy): characterization, genesis and
 226 possible hazards: *Mineralogical Magazine*, Vol. 78, pp. 919-940.

227 International Agency for Research on Cancer, Monographs, IARC, 2012, *Asbestos (chrysotile, amosite,*
 228 *crocidolite, tremolite, actinolite, and anthophyllite)*. In Evaluation of Carcinogenic Risks to Humans. A
 229 Review of Human Carcinogens. International Agency for Research on Cancer, Monographs, Vol. 100,
 230 Part C: Arsenic, Metals, Fibres, and Dusts, Lyon, France, pp. 219-309.

231 Keeling, J.L.; Raven, M.D.; and McClure, S.G., 2006, *Identification of fibrous mineral from Rowland*
 232 *Flat area, Barossa Valley, South Australia*, Primary Industries and Resources South Australia,
 233 Adelaide, Report Book 2006/00002, 23 p.

234 Leone, C., 2018, *Mineralogical and petrographic characterization of fibrous minerals and of their*
 235 *hosting rocks exposed in the north-western Voltri Group and Sestri-Voltaggio Zone, Alessandria*
 236 *province*: Unpublished post graduate thesis, Department of Earth Sciences, University of Torino, Italy,
 237 180 p.

238 Lucci, F.; Della Ventura, G.; Conte, A.; Nazzari, M.; and Scarlato, P., 2018, Naturally Occurring
 239 Asbestos (NOA) in Granitoid Rocks, A Case Study from Sardinia (Italy): *Minerals*, Vol. 8, pp. 442-
 240 464.

241 Luce, D.; Bugel, I.; Goldberg, P., Goldberg, M.; Salomon, C.; Billon-Galland, M.A.; Nicolau, J.;
 242 Quénel, P.; Fevotte, J.; and Brochard P., 2000, Environmental exposure to tremolite and respiratory
 243 cancer in New Caledonia: a case-control study: *American Journal Epidemiology*, Vol. 151, pp. 259–
 244 265.

245 Mattioli, M.; Giordani, M.; Arcangeli, P.; Valentini, L.; Boscardin, M.; Pacella, A.; and Ballirano, P.,
 246 2018, Prismatic to Asbestiform Offretite from Northern Italy: Occurrence, Morphology and Crystal-
 247 Chemistry of a New Potentially Hazardous Zeolite: *Minerals*, Vol. 8, pp. 69-84.
 248 Meeker, G.P.; Bern, A.M.; Brownfield, I.K.; Lowers, H.A.; Sutley, S.J.; Hoefen, T.M.; and Vance, J.S.,
 249 2003, The Composition and Morphology of Amphiboles from the Rainy Creek Complex, Near Libby,
 250 Montana: *American Mineralogist*, Vol. 88, pp. 1955–1969.
 251 Ministry of the Environment and Protection of the Territory and the Sea, 2018, *Asbestos mapping*:
 252 [https://www.minambiente.it/sites/default/files/bonifiche/Mappatura_amianto/mappatura_amianto_2018](https://www.minambiente.it/sites/default/files/bonifiche/Mappatura_amianto/mappatura_amianto_2018.pdf)
 253 [.pdf](https://www.minambiente.it/sites/default/files/bonifiche/Mappatura_amianto/mappatura_amianto_2018.pdf)
 254 Noonan, C. W., 2017, Environmental asbestos exposure and risk of mesothelioma: *Annals*
 255 *Translational Medicine*, Vol. 5, pp. 234-244.
 256 Paccagnella, G., 2018, *Mineralogical and petrographical characterization of asbestiform phases and of*
 257 *their host rocks from the north-western Voltri Massif*: Unpublished post graduate thesis, Department of
 258 Earth Sciences, University of Torino, Italy, 182 p.
 259 Vignaroli, G.; Ballirano, P.; Belardi, G.; and Rossetti, F., 2013, Asbestos fibre identification vs.
 260 evaluation of asbestos hazard in ophiolitic rock mélanges, a case study from the Ligurian Alps (Italy):
 261 *Environmental Earth Sciences*, Vol. 72, pp. 3679-3698.
 262 World Health Organization, WHO, 1997, *Determination of airborne fibre number concentrations; a*
 263 *recommended method, by phase contrast optical microscopy (membrane filter method)*: World Health
 264 Organization, Geneva, Switzerland, 53 p.

265 TABLE CAPTIONS

266 **Table 1.** List of NONA in decreasing order of finding frequency, detected in Piedmont and Aosta
267 Valley Regions, North-Western Alps, Italy and the ideal chemical- formula.

268 **Table 2.** Parallel intergrowths of asbestos and asbestiform minerals after investigations by TEM-EDS.
269 The most abundant asbestiform minerals (main) and the intergrown minerals (subordinate) in the same
270 bundle are listed in the left and in the right column, respectively. (asbestos are indicated in italic-
271 Italian legislation 277/91).


272


273 FIGURE CAPTIONS


274 **Figure 1.** Italian regions and mineral species of naturally occurring asbestos (NOA) in rocks:
275 chrysotile, tremolite asbestos, less diffused actinolite asbestos, and rare anthophyllite asbestos.


276 **Figure 2.** Italian regions and mineral species of naturally occurring non-asbestos classified asbestiform
277 minerals (NONA) in rocks: asbestiform antigorite, asbestiform Ca-erionite with Ca-levyne, asbestiform
278 F-edenite, asbestiform gedrite, asbestiform Mg-horneblende, asbestiform offretite, asbestiform
279 polygonal serpentine, asbestiform sepiolite. For Aosta Valley and Piedmont Regione, details are in the
280 text.**Figure 3.** Macroscopic similarity of mineral fibre bundle. A) Vein filled of asbestiform antigorite.
281 B) Fibrous bundles of asbestiform antigorite. C) Fibrous bundles of chrysotile.

282 **Figure 4.** TEM image of cross-section of an asbestiform composite made of axially-textured
283 intergrown fibrils of asbestiform antigorite (atg) with minor asbestiform polygonal serpentine (PS) and
284 chrysotile (chrys).









NONA	Ideal chemical formula
asbestiform polygonal serpentine, asbestiform antigorite	$\text{Mg}_3\text{Si}_2\text{O}_5(\text{OH})_4$
asbestiform diopside	$\text{CaMgSi}_2\text{O}_6$
asbestiform carlosturanite	$(\text{Mg,Fe,Ti})_{21}\text{Si}_{12}\text{O}_{28}(\text{OH})_4(\text{OH})_{30} \cdot \text{H}_2\text{O}$
asbestiform forsterite	Mg_2SiO_4
asbestiform balangeroite	$(\text{Mg,Fe,Mn})_{21}\text{O}_3(\text{OH})_{20}(\text{Si}_4\text{O}_{12})_2$
asbestiform sepiolite	$\text{Mg}_4\text{Si}_6\text{O}_{15}(\text{OH})_2 \cdot 4\text{H}_2\text{O}$
asbestiform talc	$\text{Mg}_3\text{Si}_4\text{O}_{10}(\text{OH})_2$
asbestiform brugnatellite	$\text{Mg}_6\text{Fe}(\text{CO}_3)(\text{OH})_{13} \cdot 4\text{H}_2\text{O}$
asbestiform brucite	$\text{Mg}(\text{OH})_2$

Table 1. List of NONA in decreasing order of finding frequency, detected in Piedmont and Aosta Valley Regions, North-Western Alps, Italy and the ideal chemical- formula.

MAIN MINERAL	SUBORDINATE MINERALS
<i>chrysotile</i> + asbestiform polygonal serpentine	asbestiform antigorite, asbestiform diopside, asbestiform carlosturanite, asbestiform forsterite, asbestiform balangeroite
asbestiform antigorite	<i>chrysotile</i> + asbestiform polygonal serpentine, <i>asbestos tremolite</i> , asbestiform carlosturanite
<i>asbestos tremolite</i> (<i>asbestos actinolite</i>)	<i>chrysotile</i> + asbestiform polygonal serpentine, asbestiform antigorite, asbestiform talc
asbestiform diopside	<i>chrysotile</i> + asbestiform polygonal serpentine, asbestiform antigorite, asbestiform carlosturanite, asbestiform balangeroite, asbestiform brugnatellite
asbestiform carlosturanite	<i>chrysotile</i> + asbestiform polygonal serpentine, asbestiform diopside, asbestiform antigorite, asbestiform forsterite, asbestiform brucite
asbestiform forsterite	<i>chrysotile</i> + asbestiform polygonal serpentine, asbestiform carlosturanite
asbestiform balangeroite	<i>chrysotile</i> + asbestiform polygonal serpentine, asbestiform diopside
asbestiform brugnatellite	asbestiform diopside
asbestiform brucite	asbestiform diopside
asbestiform sepiolite	“organic matter”

Table 2. Parallel intergrowths of asbestos and asbestiform minerals after investigations by TEM-EDS. The most abundant asbestiform minerals (main) and the intergrown minerals (subordinate) in the same bundle are listed in the left and in the right column, respectively. (asbestos are indicated in italic- Italian legislation 277/91).